

*Dispatch*

September 26, 2005 / 54(Dispatch);1-4

Infectious Disease and Dermatologic Conditions in Evacuees and Rescue Workers After Hurricane Katrina --- Multiple States, August--September,

On August 29, 2005, Hurricane Katrina struck states along the Gulf Coast of the United States. In the days after the hurricane struck, approximately 750 evacuation centers were established in at least 18 states to accommodate more than 200,000 evacuees (1). State and local health departments, with assistance from CDC, initiated infectious disease surveillance and outbreak response activities, implemented by teams of public health workers, including military personnel. Outbreak monitoring included direct reporting of conditions of public health significance to public health agencies; daily contact between CDC and local public health officials; canvassing of reports from CDC, public health departments, and news media for potential infectious disease outbreaks; and investigation of reports of infectious disease with outbreak potential. This report summarizes infectious disease and dermatologic conditions reported during the first 3 weeks after the hurricane, before effective local surveillance was fully implemented. One outbreak of norovirus was reported among evacuees in Texas; no other outbreaks requiring unusual mobilization of public health resources were reported among evacuees or rescue workers.

Dermatologic Conditions

Among hurricane evacuees from the New Orleans area, a cluster of infections with methicillin-resistant *Staphylococcus aureus* (MRSA) was reported in approximately 30 pediatric and adult patients at an evacuation facility in Dallas, Texas. Three of the MRSA infections were confirmed by culture (Figure and Table). In addition, 24 cases of hurricane-associated *Vibrio vulnificus* and *V. parahaemolyticus* wound infections (2) were reported, with six deaths.

Among rescue workers, CDC received reports of the following two types of skin lesions with infectious etiologies: tinea corporis among military personnel from two locations working in the wet environment of early evacuation efforts, and an erythematous, papular, and pustular rash consistent with folliculitis among military personnel working in Mississippi. In addition, the following three rashes subsequently determined to be noninfectious were reported in rescue workers: 1) prickly heat (miliaria crystalline, rubra, and pustulosa); 2) two clusters of nonpruritic erythematous papular, nonfollicular lesions in exposed skin of 97 military rescue workers presumed to have been caused by arthropod (likely mite) bites; and 3) circumferential lesions, appearing as macerated skin at the waist, attributed to excessive chafing.

Diarrheal Disease

CDC received reports of clusters of diarrheal disease among persons in evacuation centers in Louisiana, Mississippi, Tennessee, and Texas. In Louisiana, approximately 20 clusters of diarrheal illness in evacu-

centers were reported and investigated. In Memphis, Tennessee, gastrointestinal illness was the most common acute disease complaint among evacuees. Approximately 1,000 cases of diarrhea and vomiting were reported among adult and child evacuees in Mississippi and Texas; tests detected norovirus in stool specimens from evacuees in Texas. Sporadic nontyphoidal *Salmonella*, nontoxigenic *V. cholerae* O1, and other infections were identified. Confirmed cases of *Shigella* dysentery, typhoid fever, or infection by toxigenic *V. cholerae* O1 were reported among evacuees from Hurricane Katrina. Three weeks after the initial displacement caused by Katrina, few cases of diarrheal disease were being reported.

Respiratory Disease

Upper respiratory infections and pneumonias were reported among evacuees, including a case of pertussis in an infant aged 2 months who was rescued from a rooftop in New Orleans and evacuated to Tennessee. Appropriate antimicrobial prophylaxis was provided, and contact tracing identified no additional cases.

Control of tuberculosis (TB) among evacuees has consisted both of detecting new cases and providing continuity of care for previously known cases. A homeless person without a previous diagnosis of TB who was evacuated from New Orleans to Philadelphia was identified by entry screening with symptoms consistent with pulmonary TB. The patient was promptly isolated and began treatment for TB disease; a subsequent culture confirmed the diagnosis. At least eight other evacuees initially identified as potentially having TB were subsequently determined to have other conditions (e.g., lung cancer and infection with nontuberculous mycobacteria).

Treatment of TB requires a multidrug regimen for at least 6 months administered under directly observed therapy (DOT). A total of 195 persons in the most directly affected regions (eight counties in Alabama, six parishes in Louisiana, and 11 counties in Mississippi) were known by the local public health authorities to be under treatment for TB disease when Hurricane Katrina struck on August 29. Immediately after the hurricane, program staff sought out known TB patients to check their status and assure that therapy continued. As of September 23, all 27 currently known TB patients who resided in Alabama, all 21 in Mississippi, and 114 in Louisiana had been located through a coordinated local, state, and federal public health response. Of 41 TB patients from Louisiana not yet located, 41 were considered noncontagious at the time the hurricane struck on the basis of disease site, treatment duration, or smear status. However, treatment needs to be continued to prevent recurrence of disease and potential for emergence of drug resistance. Intense efforts continue to locate the remaining patients.

Reported by: *J Jablecki, DrPH, Div TB Control, Alabama Dept of Public Health. SA Norton, MD, 14th Support Hospital, Fort Benning, Georgia. R Keller, MD, Ochsner Clinic Foundation, New Orleans; C L. Smith, MD, TB Control Section, R Ratard, MD, S Straif-Bourgeois, PhD, Louisiana Office of Public Health; Louisiana Health and Hospitals. JM Holcombe, MPPA, S Quilter, MPH, TB Control Program; P Byers, MD, M M. Moore, MD, PhD, Mississippi Dept of Health. D Schlossberg, MD, DP Dohony, MPH, Philadelphia TB Control Program, Pennsylvania. J Neville, MD, Air Force Institute for Operational Health, Brooks City Base; J Carlo, M. Buhner, MD, Dallas County Dept of Health and Human Svcs; BR Smith, MD, C Wallace, Texas Dept of Health Svcs. D Jernigan, MD, J Sobel, MD, M Reynolds, MD, M Moore, MD, M Kuehnert, MD, J Mott, MD, Outbreak Investigations Team; G Burns-Grant, T Misselbeck, PE Cruise, P LoBue, MD, M Haddad, MSN, Div of TB Elimination, National Center for HIV, STD, and TB Prevention; TA C. Smith, Div of Bacterial and Mycotic Diseases, National Center for Infectious Diseases; A Cohen, MD, R Sune, MD, M Jhung, MD, P Vranken, DPH, FMT Lewis, MD, LR Carpenter, DVM, EIS officers, CDC.*

Editorial Note:

Environmental conditions after natural disasters increase the risk for infectious disease. The experience of the public health community with diarrheal disease after natural disasters suggests that evacuation centers s

prepared for considerable demands on clinic staff, janitorial services, and maintenance of personal hygiene. Functioning flush toilets and potable water are often impaired. Extensive flooding, such as that which Katrina, can increase risk for exposure to waterborne agents and vectors such as mosquitoes (4). Hurricane survivors can suffer wound injuries that can become infected. Persons unable to evacuate affected areas underlying illnesses might be more susceptible to infectious disease. Finally, once evacuees have relocated to evacuation centers, crowding and unsanitary conditions can amplify transmission of infectious disease. Despite the massive migration of evacuees and their subsequent placement in evacuation centers, only one outbreak of communicable disease (norovirus) requiring unusual mobilization of public health resources was reported as of September 23.

Observations from previous natural disasters, such as the 2004 Asian tsunami (5), suggest that skin, diarrheal, and respiratory infections are the most common infectious diseases in survivors. Infectious disease outbreaks are rare following natural disasters, especially in developed countries (6,7), and specific etiologies are unpredictable, reflecting infectious diseases endemic to the affected region before the disaster. Injury and infections are expected during the first few days after a disaster. In contrast, airborne, waterborne, and foodborne diseases are expected to occur for up to 1 month after a disaster. Because norovirus is a common cause of illness that is easily spread in densely populated communities (e.g., classrooms and cruise ships), outbreaks are anticipated after a natural disaster. Conversely, the rarity of toxigenic *V. cholerae*, dengue, and malaria in the Gulf Coast states before Hurricane Katrina suggested that they were unlikely to pose a public health challenge after the hurricane. Another consequence of natural disasters is the potential exposure to dead bodies, both human and animal. No evidence exists that exposure to bodies after a disaster leads to infectious disease epidemics. Persons handling human corpses and animal carcasses might be exposed to infectious pathogens and should use appropriate protective equipment (8).

Hurricane Katrina affected an area of approximately 90,000 square miles and resulted in the displacement of approximately 1 million persons. The local and state public health infrastructure for communicable disease surveillance and control was disrupted, shifting initial emphasis toward nonstandardized mechanisms for reporting. These mechanisms included direct reporting to public health agencies, which resulted in timely recognition of suspected instances of infectious disease while effective local surveillance was being established. The main offices of the Louisiana Office of Public Health were affected by the hurricane, and all operations were temporarily relocated from New Orleans to Baton Rouge. Because of the compromised capacity of the Louisiana State Public Health Laboratory, laboratory confirmation of causative agents was relocated to regional public health laboratories. Effective surveillance is now under way in all of the affected states.

Guidance for health-care professionals for preventing and responding to potential infectious disease outbreaks after a hurricane is available at <http://www.bt.cdc.gov/disasters/hurricanes/hcp.asp>. In addition to routine vaccinations, evacuees living in crowded groups should consider vaccinations for influenza, varicella, and mumps-rubella (MMR), and hepatitis A (available at <http://www.bt.cdc.gov/disasters/hurricanes/immunizations.asp>). However, immunocompromised persons, persons infected with human immunodeficiency virus, pregnant women, and persons on systemic steroids should not receive the live viral vaccines, varicella and MMR. Screening should be performed by self-report.

Methods used by TB outreach workers to locate persons with previous TB infections include 1) asking whether they have heard from patients since the hurricane, 2) visiting locations known to be frequented before the hurricane, 3) calling all known phone numbers for each patient, 4) checking public registries such as the American Red Cross Family Linking at <http://www.katrinSAFE.org> for updated news on patient whereabouts, 5) seeking access to mortality records and other registries for cross-matching. A Katrina Help Desk was established on September 2 at the CDC Division of Tuberculosis Elimination to facilitate communication between the National TB Controllers Association and state TB programs as patients were relocated to other state care providers should immediately contact their state TB controllers (available at <http://www.cdc.gov/nchstp/tb/katrina/tbcontrollers.htm>) if an evacuee has TB symptoms (i.e., cough of

duration, fever, chills, night sweats, or recent weight loss) or was under treatment for TB disease. The I TB Control Program (telephone 337-262-5616) will provide information about treatment regimens previously administered to relocated patients. If providers are unable to reach a local or state TB controller, they may contact the CDC Division of Tuberculosis Elimination at telephone 404-639-8336 for assistance.

Acknowledgments

This report is based, in part, on contributions by NB Keenon, MPH, K Taylor, MPH, Div of TB Control, Louisiana Dept of Public Health. G Barre, MD, Ochsner Clinic Foundation, New Orleans; J Plough, Louisiana Dept of Public Health; Louisiana Dept of Health and Hospitals. B Amy, MD, S Hand, Mississippi Dept of Health and Hospitals, Children's Medical Center of Dallas; J Romero, MPH, Dallas County Dept of Health and Human Services; M Duran, MD, PhD, A Groda, AJ Abell, PhD, MA Davis, MS, Texas Dept of State Health Services.

References

1. CDC. Update on CDC's response to Hurricane Katrina: from the CDC Director's Emergency Operations Center, p.m. update, September 6, 2005. Available at <http://www.cdc.gov/od/katrina/09-06-05.htm>
2. [CDC. *Vibrio* illnesses after Hurricane Katrina---multiple states, August--September 2005. MMWR 2005;54:928--31.](#)
3. [CDC. Treatment of tuberculosis: American Thoracic Society, CDC, and Infectious Diseases Society of America. MMWR 2003;52\(No. RR-11\).](#)
4. [CDC. Tropical Storm Allison rapid needs assessment---Houston, Texas, June 2001. MMWR 2001;50:9.](#)
5. Lim JH, Yoon D, Jung G, Joo Kim W, Lee HC. Medical needs of tsunami disaster refugee camps. *Emerg Infect Dis* 2005;37:422--8.
6. Shultz JM, Russell J, Espinel Z. Epidemiology of tropical cyclones: the dynamics of disaster, disaster development. *Epidemiol Rev* 2005;27:21--35.
7. Blake PA. Communicable disease control [Chapter 3]. In: *The public health consequences of disasters*. 1989. MB Gregg, ed. Atlanta, GA: US Department of Health and Human Services, CDC; 1989.
8. Morgan O. Infectious disease risks from dead bodies following natural disasters. *Rev Panam Salud Publica* 2004;15:307--12.

Figure

FIGURE. Methicillin-resistant *Staphylococcus aureus* in the leg of an evacuee from Hurricane Katrina — Dallas, Texas, September 2005



Photo/P Hicks, Children's Medical Center of Dallas

[Return to top.](#)

Table

TABLE. Number of cases of selected diseases and conditions reported in evacuees and rescue workers during the 3 weeks immediately after Hurricane Katrina made landfall — multiple states, August–September 2005

Disease/Condition*	No. of cases	States reporting	
Dermatologic conditions			
<i>Infectious</i>			
Methicillin-resistant <i>Staphylococcus aureus</i> infections	30 (3 confirmed)	Texas	Ev
<i>Vibrio vulnificus</i> and <i>V. parahaemolyticus</i> wound infections	24 (6 deaths)	Arkansas, Arizona, Georgia, Louisiana, Mississippi, Oklahoma, Texas	Ev
Tinea corporis	17	Mississippi	Re
<i>Noninfectious</i>			
Arthropod bites (likely mite)	97	Louisiana	Re
Diarrheal disease			
Acute gastroenteritis, some attributed to norovirus	Approximately 1,000	Louisiana, Mississippi, Tennessee, Texas	Ev
Nontoxigenic <i>V. cholerae</i> O1	6	Arizona, Georgia, Mississippi, Oklahoma, Tennessee	Ev
Nontyphoidal <i>Salmonella</i>	1	Mississippi	Ev
Respiratory disease			
Pertussis	1	Tennessee	Ev
Respiratory syncytial virus	1	Texas	Ev
Streptococcal pharyngitis	1	Texas	Ev
Tuberculosis	1	Pennsylvania	Ev
Other condition			
Presumed viral conjunctivitis	Approximately 200	Louisiana	Ev

* Other diseases and conditions, for which the number of cases was unknown, included scabies; circumferential lesions at waist; contact dermatitis; papular, pustular rash consistent with folliculitis; immersion foot; prickly heat; influenza-like illness and upper respiratory infections; and head li

[Return to top.](#)

Use of trade names and commercial sources is for identification only and does not imply endorsement by the U.S. Department of Health and Human Services.

References to non-CDC sites on the Internet are provided as a service to *MMWR* readers and do not constitute or imply endorsement of these organizations or their programs by CDC or the U.S. Department of Health and Human Services. CDC is not responsible for content of pages found at these sites. URL addresses listed in *MMWR* were current as of the date of publication.

Disclaimer All *MMWR* HTML versions of articles are electronic conversions from ASCII text into HTML. This conversion have resulted in character translation or format errors in the HTML version. Users should not rely on this HTML document, referred to the electronic PDF version and/or the original *MMWR* paper copy for the official text, figures, and tables. An original copy of this issue can be obtained from the Superintendent of Documents, U.S. Government Printing Office (GPO), Washington 20402-9371; telephone: (202) 512-1800. Contact GPO for current prices.

**Questions or messages regarding errors in formatting should be addressed to mmwrq@cdc.gov.

Date last reviewed: 9/26/2005

[HOME](#) | [ABOUT *MMWR*](#) | [MMWR SEARCH](#) | [DOWNLOADS](#) | [CONTACT](#)
[POLICY](#) | [DISCLAIMER](#) | [ACCESSIBILITY](#)

SAFER • HEALTHIER • PEOPLE™

Morbidity and Mortality Weekly Report
Centers for Disease Control and Prevention
1600 Clifton Rd, MailStop K-95, Atlanta, GA 30333,
U.S.A



Department of Health
and Human Services